



Name :

Roll No. :

Invigilator's Signature :

CS/MCA/SEM-1/M(MCA)-101/2012-13

2012

DISCRETE MATHEMATICAL STRUCTURES

Time Allotted : 3 Hours

Full Marks : 70

The figures in the margin indicate full marks.

Candidates are required to give their answers in their own words as far as practicable.

GROUP – A

(Multiple Choice Type Questions)

1. Choose the correct alternatives for any *ten* of the following :

10 × 1 = 10

i) Out of the following the singleton set is


a) $A = \{ x : 3x - 2 = 0, x \in Q \}$

b) $B = \{ x : x^2 - 1 = 0, x \in R \}$


c) $C = \{ x : 30x - 59 = 0, x \in N \}$

d) $D = \{ x : x^2 - 1 = 0, x \in Z \}$

where Q, R, N, Z is the set of all rational number, real number, natural number and integers respectively.



- ii) If A, B & C are any three arbitrary sets, then $A - (B \cap C)$ is
- a) $(A - B) \cup (A - C)$ b) $(A - B) \cap (A - C)$
 c) $(A - B) \cap (C - A)$ d) $(B - A) \cup (A - C)$.
- iii) The number of arrangements of 25 objects where 7 are of the first kind, 2 are of the second kind, 3 are of the third kind and 4 are of the fourth kind is given by
- a) $\frac{25!}{7!2!3!4!}$ b) $\frac{25!}{7!2!}$
 c) $\frac{25!}{3!4!}$ d) none of these.
- iv) Out of the following statements the formula for tautology is
- a) $(p \vee q) \rightarrow q$ b) $p \vee (q \rightarrow p)$
 c) $p \vee (p \rightarrow q)$ d) $p \rightarrow (p \rightarrow q)$.
- v) The solution of the recurrence relation $a_r - 7a_{r-1} + 10a_{r-2} = 0$ given $a_0 = 0, a_1 = 3$ is
- a) $5^{-r} - 2^r$ b) $5^r + 2^r$
 c) $5^r - 2^r$ d) none of these.
- vi) The type of the grammar, which consists of the productions $s \rightarrow aA, A \rightarrow aAB, B \rightarrow b, A \rightarrow a$ is
- a) Type-0 b) Type-1
 c) Type-2 d) Type-3.



vii) Let L be a language given by $L = \{a^n b^n : n \geq 0\}$, then L^2 is equal to

a) $\{a^n b^n a^m b^m : n \geq 0, m \geq 0\}$

b) $\{a^n b^n : n \geq 0\}$

c) $\{a^n b^n a^n b^n : n \geq 0\}$

d) none of these.

viii) The coefficient of X^{25} in $(X^3 + X^4 + X^5 + \dots)^5$ is

a) $C(9, 5)$

b) $C(5, 9)$

c) $C(5, 5)$

d) $C(9, 9)$.

ix) For the mapping $g : [-3, 2] \rightarrow R$ defined by $g(x) = 3x + 4$ for any $x \in [-3, 2]$ then image set of g is

a) $[-5, 10]$

b) $[0, 10]$

c) $[2, -3]$

d) none of these.

x) A spanning tree of a connected graph contains

a) all the vertices of the graph

b) all the vertices and edges of the graph

c) a few vertices of the graph

d) none of these.

xi) If a binary tree has 20 pendant vertices then the number of internal vertices of the tree is

a) 20


b) 21

c) 23

d) 19.



xii) Haase diagram is given below :



This is a


- a) Poset
- b) Toset
- c) Lattice
- d) none of these.

GROUP - B


(Short Answer Type Questions)

Answer any *three* of the following. $3 \times 5 = 15$

2. Define adjacency matrix of a simple graph $G = (V, E)$. Write down the adjacency matrix for the following undirected graph :



3. By using Principle of Mathematical Induction, prove that $4^{2n+1} + 3^{n+2}$ is an integer multiple of 13 for all positive integers n .
4. Let $A = \{x \in R : x \neq 2\}$ & $B = \{x \in R : x \neq 1\}$, and let the two functions $f : A \rightarrow B$ & $g : B \rightarrow A$ are defined by $f(x) = \frac{x}{x-2}, \forall x \in A$ and $g(x) = \frac{2x}{x-1}, \forall x \in B$, then find $f \circ g$. Are the two functions f and g invertible ? 2 + 3



5. Over the alphabet $\Sigma = \{a, b\}$ design a DFA which accepts the language $L = \{w : w \text{ has both an even number of } a\text{'s and an even number of } b\text{'s}\}$.
6. Find an explicit formula for the sequence defined by $a_n = a_{n-1} + 4 \quad \forall n \geq 2$ with $a_1 = 2 \dots$.

GROUP - C

(Long Answer Type Questions)

Answer any *three* of the following. $3 \times 15 = 45$

7. a) Determine the intersection of the following two fuzzy sets :



$$A = \left\{ \frac{4}{0.1}, \frac{6}{0.5}, \frac{8}{0.6}, \frac{10}{0.7} \right\} \text{ and}$$

$$B = \left\{ \frac{0}{0.4}, \frac{2}{0.6}, \frac{4}{1}, \frac{6}{1}, \frac{8}{0.6}, \frac{10}{0.5} \right\}.$$

- b) For each of the following mappings determine whether it is (i) injective, (ii) surjective. Find the inverse mapping of the mapping which is bijective.

$$K : \mathbb{R} \rightarrow \mathbb{R} \text{ defined by } k(x) = \begin{cases} x^2 - 1, & x \geq 0 \\ -x^2 - 1, & x < 0 \end{cases}$$

- c) Examine if the following graphs are isomorphic :



3 + 7 + 5

8. a) Solve the following recurrence relation using generating function :

$$a_n - 9a_{n-1} + 20a_{n-2} = 0 \text{ for } n \geq 2 \text{ and } a_0 = -3, a_1 = -10.$$



- b) Show that $n^2 > 2n + 1$ for $n \geq 3$ using mathematical induction.
- c) Show that $(p \vee q) \wedge (\neg p \wedge \neg q)$ is a contradiction.

7 + 4 + 4

9. a) Write DNF of the following statement :

$$\neg \{ \neg (p \leftrightarrow q) \wedge r \}$$


- b) Verify whether the argument given below is valid or not :
All mammals are animals. Some mammals are two-legged. Therefore, some animals are two-legged.

- c) Prove the following equivalence :

$$\neg p \wedge q \Leftrightarrow \neg (p \vee (\neg p \wedge q))$$

5 + 5 + 5

10. a) Find by Prim's algorithm a spanning tree with minimum weight from the graph given below. Also calculate total weight of spanning tree :



- b) Prove that a connected graph n with $n - 1$ vertices and edges is a tree.
- c) Determine the value of n if $4 \times {}^n P_3 = {}^{n+1} P_3$. 6 + 6 + 3

11. a) Prove that in a bounded distributive lattice (L, \cap, \cup) an element cannot have more than one complement.
- b) Find the sum of all four digits for even numbers that can be made with the digits 0, 1, 2, 3, 5, 6 & 8.



- c) Define Mealy and Moore machine. Construct a Moore machine from the following Mealy machine :

Present State	Next State			
	$a = 0$		$a = 1$	
	State	Output	State	Output
s_0	s_0	1	s_1	0
s_1	s_3	1	s_3	1
s_2	s_1	1	s_4	1
s_3	s_2	0	s_0	1

4 + 6 + 5

=====